

Supporting Information

Manuscript Title:

Projection of corn production and stover-harvesting impacts on soil organic dynamics in the U.S. Temperate Prairies

Authors:

Yiping Wu^{1*}, Shuguang Liu^{2*}, Claudia J. Young³, Devendra Dahal⁴, Terry L. Sohl², Brian Davis⁴

¹ ASRC Federal, contractor to U.S. Geological Survey (USGS) Earth Resources Observation and Science (EROS) Center, Sioux Falls, SD 57198, USA

Email: ywu@usgs.gov (Y. Wu)

² U.S. Geological Survey (USGS) Earth Resources Observation and Science (EROS) Center, Sioux Falls, SD 57198, USA

Email: slu@usgs.gov (S. Liu)

³ Innovate! Inc., contractor to the USGS EROS Center, Sioux Falls, SD 57198

⁴ Stinger Ghaffarian Technologies, contractor to the USGS EROS Center, Sioux Falls, SD 57198

Projected land cover trend

Based on the FORE-SCE land cover model projections, the annual proportions of land-cover classes during 2001 through 2050 under the four IPCC scenarios (A1B, A2, B1, and B2) are presented below.





Figure S.1 Annual time series of land-cover class proportions in the Temperate Prairies region from 2001 to 2050 under the four scenarios (A1B, A2, B1, and B2).

31 **Location of the study area**



32

33 [Figure S.2](#) Location and the land cover and land use map (2001) of the Temperate

34 Prairies (Ecoregion 9.2). The map was created using ArcGIS10.2.

35